

P-26

Diversity of *Fusarium proliferatum* Isolated from Asparagus Plants in Reference to Virulence, Toxicogenicity and Patterns of DNA-Analysis

W. Xu¹, A. Kofoet¹, M. Gossmann² and H.-U. Humpf³

¹ Institute for Vegetable and Ornamental Crops, Grossbeeren/Erfurt e. V., 14979 Grossbeeren, Germany

² Institute of Horticultural Science, Humboldt University of Berlin, 14195 Berlin, Germany

³ Institute of Food Chemistry, University of Muenster, 48149 Muenster, Germany

Fusarium proliferatum was detected frequently as one of the dominant *Fusarium* species in asparagus rhizome. Isolates of this species were obtained from both symptomatic and non-symptomatic plants. Virulence tests and toxicogenicity analysis regarding fumonisin production were carried out in this present study to assess the relevant characteristics of the populations of *F. proliferatum* infecting plants concerning both plant health and food safety. The results showed (1) The isolates both from symptomatic and non-symptomatic plants were differently virulent; (2) Isolates from symptomatic plants could be less virulent than those from non-symptomatic plants and vice versa; (3) Isolates from the same plant could be differently virulent; (4) All tested isolated could produce fumonisins in vitro, (5) but the abilities of fumonisin production differed quantitatively among the isolates; (6) No correlation between virulence and toxicogenicity could be demonstrated in our tests. In addition to the tests above, DNA of the isolates of *F. proliferatum* were investigated by use of RAPD-PCR to determine the genetic diversity of the isolates. According to the patterns of the DNA-analysis the isolates fell into different groups, but a correlation of this grouping to virulence or toxicogenicity of the respective isolates is unclear as yet and needs further investigation. Additionally, the results showed that isolates within a plant were genetically close to each other, but must not absolutely identical.