Detection of a new ilarvirus in Passiflora edulis

Christian Lüchau¹, Joseph Cutler¹, Juliane Langer¹, Susanne von Bargen¹, Orlando Acosta Losada², Gerhard Fischer², Fánor Casierra-Posada³, Adriana Castañeda Cárdenas⁴, Mónica Betancourt Vasquez⁵, Wilmer Cuellar⁶, Eduardo Arvydas Stasiukynas⁷, Emilio Arevalo-Peñaranda⁴, Carmen Büttner¹*

¹ Humboldt Universität zu Berlin, Germany ² Universidad Nacional Bogotá, Colombia ³ Universidad Pedagógica y Tecnológica Tunja, Colombia ⁴ Instituto Colombiano Agropecuario, Bogotá, Colombia, ⁶ International Center for Tropical Agriculture (CIAT) Cali, Colombia ⁷ Hacienda Misiones, Cundinamarca, Colombia ⁵ Corporación Colombiana de Investigación Agropecuaria, Bogotá, Colombia, ⁶ International Center for Tropical Agriculture (CIAT) Cali, Colombia ⁷ Hacienda Misiones, Cundinamarca, Colombia

*phytomedizin@agrar.hu-berlin.de



Introduction

Colombia is one of the world's main producers and exporters of tropical fruits, however, it lacks a robust preventive management programme for the control of plant viruses¹.

Recent Next Generation Sequencing (NGS) according to Massart et al. (2017)² have shown the presence of a new ilarvirus in Colombian purple passion fruit farms. The spread of this virus could mean drastic reduction of crop yields and major economic losses. For this purpose, molecular biological tools such as nucleic acid isolation, polymerase chain reaction (PCR), and NGS are being used to measure the frequency and distribution of this virus in Colombia, to characterise which symptoms are associated with it, and to identify the pathways for its transmission. This research will contribute to methods for distributing virus-free plant material of Passiflora edulis in Colombia.













Figure 1. Maps of Colombia, where the new ilarvirus in purple passion fruit was found, and the region where the samples were taken from (a-b). Purple passionfruit field in Mesitas del Colegio, Colombia (c).

Materials and Methods

• Samples of *P. edulis* plant material with virus-suspected symptoms were collected in 2016 and 2017 from 5 different farms in the Altiplano Cundiboyasence Andean plateau (Cundinamarca and Boyaca departments, Colombia). 41 samples from 2 of these 5 farms, were used in this study.

•NGS analysis revealed 57% identity to the movement protein (MP) of Lilac ring mottle virus (*llarvirus*), and 65% identity to the replicase of the *Tomato* necrotic streak virus (llarvirus)

• RNA isolation according to Boom et al. (1990)³ from leaf material

- cDNA synthesis with pMMLV RTase (fresh material) or Premium RevertAid RTase (frozen material) and random hexamer primers
- •nad5-PCR for quality control of RNA and cDNA synthesis according to Menzel et al. $(2002)^4$
- Diagnostic RT-PCR for the detection of the novel ilarvirus with primers developed for its RNA1, RNA2, and RNA3
- •RT-PCR for viral-specific detection of the CP coding region of RNA3 of the novel ilarvirus



Figure 1. Genome structure of the *llarvirus* genus, with a tripartite linear ssRNA(+) genome. The positions of

Figure 3. Detection of the new ilarvirus by means of RT-PCR, using designed primers based on the previously obtained NGS analysis. RNA1, RNA2 and RNA3 diagnostic PCRs, and primers targeting RNA3 N-terminal region, RNA3 Cterminal region, and RNA3 complete CP Region. In all PCRs, one positive control (+), previously tested, and water or a healthy control, were used as negative controls (-). Numbers from 1-19 refer to the same samples that were tested in the different PCRs.

Table 1. Results from the detection of the new ilarvirus by means of diagnostic and genus-specific RT-PCRs, using RNA1, RNA2, and RNA1 designed primers. Total number of samples processed: 41.

Symptom	Blistering	Yellowing	Chlorosis	Blistering/ Deformation	Blistering/ Chlorosis	Deformation/ Chlorosis	Healthy
Samples with symptom	5	2	8	12	7	4	2
RNA1	2	0	1	4	2	0	0
RNA2	2	0	0	4	2	0	0
RNA3 Diagnostic	2	0	1	4	2	0	0
RNA3 N- terminal Region	2	0	1	4	2	0	0
RNA3 C- terminal Region	2	0	0	3	2	0	0
RNA3 Complete CP Region	2	0	0	3	0	0	0

the developed primers are shown by blue arrows. For the RNA3 primers the different combinations of the CP region are shown (a). Physical morphology of the ilarviruses (b). Figure adapted from ViralZone (2009).

Conclusions

The new ilarvirus found in the purple passion fruit (*Passiflora edulis* Sims) in Colombia can be detected by RT-PCR, using different types of primer combinations of the 3 RNAs of its genome. Therefore, this could be a useful tool to detect the presence of the virus in the passion fruit fields.

As seen in Table 1., the virus could be detected in samples showing blistering, blistering/deformation, and blistering/chlorosis symptoms. The coat protein (CP) gene is generally more variable, and therefore, characterisation of the virus, is best done by amplification of the N-terminal and C-terminal regions of the CP region by separate RT-PCRs.

References

Special thanks to our partners for diagnostic and logistical support:

¹Rodríguez, M. H., Niño, N. E., Cutler, J., Langer, J., Casierra, F., Miranda, D., Bandte, M., Büttner, C. (2016). Certificación de material vegetal sano en Colombia: un análisis crítico de oportunidades y retos para controlar enfermedades ocasionadas por virus. Revista Colombiana de Ciencias Hortícolas 10(1), pp. 164-175.

²Massart, S., Candresse, T., Gil, J., Lacomme, C., Predajna, L., Ravnikar, M., Reynard, J., Rumbou, A., Saldarelli, P., Skoric, D., Vainio, E., Valkonen, J., Vanderschuren, H., Varveri, C., Wetzel, T. (2017). A Framework for the Evaluation of Biosecurity, Commercial, Regulatory, and Scientific Impacts of Plant Viruses and Viroids Identified by NGS Technologies. Frontiers in Microbiology, 8, 45.

³Boom, R., Sol, C. J. A., Salimans, M. M. M., Jansen, C. L., Wertheim-van Dillen, P. M. E., van der Noordaa, J. (1990). Rapid and Simple Method for Purification of Nucleic Acids. Journal of Clinical Microbiology, 28 (3), pp. 495-503.

⁴Menzel, W., Jelkmann, W., Maiss, E. (2002). Journal of Virological Methods, 99, pp. 81-92.

